



## Weight-Length Models and Relative Condition Factors of Nine (9) freshwater fish species from the Yapei Stretch of the White Volta, Ghana

Seth Mensah Abobi

Department of Fisheries and Aquatic Resources Management, University for Development Studies, P.O. Box TL 1882 Tamale, Ghana.

### ARTICLE INFO

#### Article history:

Received: 4 November 2014;

Received in revised form:

20 January 2015;

Accepted: 12 February 2015;

#### Keywords

Weight-length relationship,

Condition factor,

Freshwater,

White Volta.

### ABSTRACT

Weight-Length relationships (WLRs) and Relative condition factors (Kn) were estimated for 9 freshwater fish species namely *Brycinus leuciscus* (Günther, 1867), *Hydrocynus forskali* (Cuvier, 1819), *Odaxothissa mento* (Regan, 1917), *Labeo coubie* (Rüppell, 1832), *Labeo parvus* (Boulenger, 1902), *Labeo senegalensis* (Valenciennes, 1842), *Distichodus enycephalus* (Günther, 1864), *Parailia pellucida* (Boulenger, 1901) and *Schilbe mystus* (Linnaeus, 1758) from the lower reaches of the White Volta. Samples were collected from the catches of the artisanal fishermen from October 2011 to March 2012. Using  $W=aTL^b$ , the weight-length relationships of the species were calculated. Relative condition factors of the species were obtained using the formula:  $W_o/W_c$ . The WLRs of all the 9 fish species were highly significant ( $P<0.001$ ). The determination coefficients ( $r^2$ ) ranged between 0.5764 for *Parailia pellucida* to 0.9784 for *Hydrocynus forskali*. The slopes ( $b$  values) of the WLR regression models of the nine (9) species ranged from 2.0045 to 3.1614 with 95% confidence interval of  $b$  for all the nine fish species within 1.386 to 3.352. In terms of growth pattern, 4 out of the 9 species exhibited isometric growth ( $b=3$ ). Three (3) were observed to have negative allometric growth ( $b<3$ ) and 2 fish species had positive allometric growth ( $b>3$ ) patterns. Mean relative condition factors of the species ranged from 1.00 to 1.09. A T-test of the observed weight ( $W_o$ ) versus the calculated ( $W_c$ ) on all the nine (9) species produced no significant difference ( $P>0.05$ ) between the observed weight and the calculated weight. The models are thus good proxies for approximating the weights of the species. Even though, not all the species exhibited isometric growth, the general wellbeing of the fish species was suitable during the study period as inferred from relative condition factors (Kn). This is a contribution of information to support researchers in future and fishery managers for reliable growth estimation and health status of freshwater fishes.

© 2015 Elixir All rights reserved.

### Introduction

Weight-length relationships (WLRs) and condition factor are of great importance to the fishery industry as they help to predict the best length and time suited to harvest a particular species of fish. The estimation of population size of a fish stock for the purpose of its rational exploitation often requires knowledge of individual body WLRs in the population (Dulčić and Kraljević, 1996). WLRs have several applications, namely on fish biology, physiology, ecology and fisheries assessment. In biological studies, WLR enables seasonal variation in fish growth to be followed and the calculation of condition indices. WLR gives us life history and morphological comparisons between different fish species or between different fish populations from different habitats (Gonçalves et al., 1997; Petrakis and Stergiou, 1995; Santos et al., 2002). This study was conducted to determine WLRs of 9 fish species from the lower reaches of the White Volta at Yapei, Northern Region of Ghana.

According to Abobi and Ekau (2013), the importance of the Yapei fishery cannot be underestimated. Yapei is 52 km from the Northern regional capital Tamale. Tamale is the third largest city in Ghana with a population of 537,986 (Wikipedia, 2012). Yapei is easily accessible as it is on the Accra-Kumasi-Tamale trunk road. Yapei is one of the 9 main sources of smoked fish to

the Tamale Central Market. The Tamale Central Market is the largest market center in the metropolis. It thus makes Yapei an important supplier of fish especially smoked fish. Fishermen at the Yapei stretch of the White Volta River practice unregulated and unselective harvesting of fish with fishermen harvesting all sizes of fish without regard to the sustenance of the fishery. These practices targeted at more catches for more income. Fish can only be harvested at the maximum sustainable yield (MSY) when all the biological parameters are known (Abobi and Ekau 2013).

Yapei is thus the terminal point for the lake Volta and the lower reaches of the White Volta River (WVR) starts from there. According to Braimah (2003) some 300 000 fisher folk depend on the Volta reservoir for their livelihood. The Volta reservoir is endowed with fisheries resources. Dankwa et al. (1999) identified 121 species in the Volta reservoir.

The Lake Volta and its tributaries in the last four decades have undergone great changes in its ecology, limno-chemistry and socio-economy. Increased pressure on land along the banks has led to high rates of deforestation. This has resulted in increased soil erosion leading to the transportation of high loads of silt and nutrients through rivers into the lake, there by

Tele:

E-mail addresses: [mabobi@uds.edu.gh](mailto:mabobi@uds.edu.gh)


contributing to its eutrophication. Furthermore, wetlands bordering the lake are being converted into agricultural land or land for grazing cattle, and therefore may not be able to act as natural filters for nutrients and silt, and now do not provide breeding grounds for many fish species (Ofori-Danson *et al.*, 2001). As a result of these problems, the Lake Volta Research and Development Project (VLR&DP) was undertaken under the Food and Agricultural Organization and United Nations Development Programme during the first decade of the lake's existence (FAO and UNDP 1971, 1979). These studies came to an end in 1978. Since then, systematic data collection from the Lake Volta and Volta Rivers natural resources has been lacking. There have been calls for renewed studies to facilitate their management due to declining catches.

Understanding the weight-length relationship is of paramount importance in fishery resource management and useful in comparing life history and morphological aspects of populations inhabiting in different regions (Karna *et al.*, 2012). Furthermore, this relationship is used to ascertain the condition of fish and to determine whether the growth pattern is isometric or allometric. Besides, the weight-length measurement also provides information on the stock composition, life span, mortality, growth and production of fish (Bolgar and Connolly, 1989; King, 1996; Moutopoulos and Stergiou, 2000). The relative condition factor is used to know the variation between observed and expected weight of fishes (Kund *et al.*, 2011). Therefore, in the present study, an attempt has been made to estimate the weight-length relationships and relative condition factors of *Brycinus leuciscus*, *Hydrocynus forskali*, *Odaxothrissa mento*, *Labeo coubie*, *Labeo parvus*, *Labeo senegalensis*, *Distichodus engycephalus*, *Parailia pellucida* and *Schilbe mystus* harvested from the lower reaches of the White Volta, Ghana. Such information is useful in understanding the key biological attributes of these species and their sustainable management.

## Materials and Methods

### Study area

The samples were conducted at the Yapei stretch of the White Volta River (Figure 1), which is a major landing site in the Northern Region of Ghana. Yapei is located within the Central Gonja District of the Northern Region of Ghana. It lies within latitudes 9° 10' 0" N, 1° 10' 0" W.



**Figure 1. Map of Ghana with an insert showing Central Gonja District and study area Yapei**


### Fish sampling, measurement and analysis

The samples of the selected fish species were taken at random from landings and their total length recorded to the nearest 0.1 cm using a fish measuring board while the body weight was recorded to the nearest 0.01 g using a balance after using tissue to mop off water from the surface of the specimens. The length-weight equation;  $W = aTL^b$  as described by Ricker (1975) was used to establish the length-weight relationship of the species measured, where  $W$  is the weight (g),  $TL$  is the total

length (cm),  $a$  is a constant determined empirically and  $b$  is the slope of the equation. The degree of relationship between the variables was computed by the determination coefficient ( $r^2$ ). The slope  $b$  has a numerical value which is mostly between 2.5 and 3.5 and is often close to 3 (Bagenal and Tesch, 1978; Carlander, 1969; Gayanilo and Pauly, 1997; Froese, 2006; Weatherley and Gill, 1987). Under isometric growth  $b$  equals to 3 but isometric growth in fish is rare (Bolger and Connolly, 1989; McGurk, 1985). The 95% confidence limits for  $b$  (CL 95%) were calculated using the equation:  $CL = b \pm (1.96 \times SE)$  where  $SE$  is the standard error of  $b$ . The values of the exponent  $b$  provide biological information on kind/pattern of growth of fish. The growth is isometric if  $b=3$  and the growth is allometric if  $b \neq 3$  (negative allometric if  $b < 3$  and positive allometric if  $b > 3$ ). In order to check if the value of  $b$  was significantly different from 3, the Student's  $t$ -test was conducted as expressed by the equation according to Sokal & Rohlf (1987):  $ts = (b-3)/SE$ , where  $ts$  is the  $t$ -test value,  $b$  the slope and  $SE$  the standard error of the slope  $b$ . The test statistics ( $ts$ ) values were compared with critical values from the T-Table. All the statistical analyses were considered at significance level of 5% ( $p < 0.05$ ).

The equation relating weight-length gives some indication of the state of wellbeing of a fish in a population which is known as the condition factor. Fish in good condition have higher condition factor value than those in poor condition. The relative condition factor ( $K_n$ ) which is a ratio of observed weight ( $W_o$ ) and calculated weight ( $W_c$ ) that is  $W_o/W_c$  was estimated as suggested by Le Cren (1951). The relative condition factor being one or closer to one indicates, the good condition of fish.


All calculations were performed using descriptive statistics of the Software STATISICA 8.1 and Microsoft Excel 2013. They were used to analyse weight-length data for means, deviations, frequencies, percentages, weight-length power regression functions, relative condition factors and T-test of significant difference between observed weight and calculated weight.




**Figure 2. Length-weight relationships *Hydrocynus forskali*, *Brycinus leuciscus* and *Odaxothrissa mento* from the Yapei stretch of the White Volta River. N, Number of fish; W, Weight (grams) and TL, Total Length (cm)**

## Results and Discussion

A total of 741 individuals belonging to 9 species from 5 families were sampled for this study. Cyprinidae had 3 species, Alestidae and Schilbedae were represented by two species each while Clupeidae and Distichodontidae had one species each. The parameters of the weight-length relationships of *Brycinus leuciscus*, *Hydrocynus forskali*, *Odaxothrissa mento*, *Labeo leuciscus*, *Labeo parvus*, *Labeo senegalensis*, *Distichodus engycephalus*, *Parailia pellucida* and *Schilbe mystus* are presented in Table 1 with their scatterplots in Figures 2-4. The weight-length relationships (WLRs) were highly significant ( $P < 0.001$ ) for all the 9 fish species. The determination coefficients ( $r^2$ ) ranged between 0.5764 for *Parailia pellucida* to 0.9784 for *Hydrocynus forskali*. The estimates of  $b$  ranged from 2.0045 in *Parailia pellucida* to 3.1614 in *Labeo coubie* with 95% confidence interval of  $b$  for all the nine fish species ranging from 1.386 to 3.352. The finding on the  $b$  estimates of this study is similar to the observations of Enstua-Mensah *et al.* (1995) who reported that estimates of  $b$  ranged from 2.32 to 3.27 for 45 fish species (which included species from all the families of this study) sampled in the Oti, Pru and Black Volta rivers of Ghana. However, confidence interval of  $b$  for all the nine fish species of this present study is different from confidence interval of  $b$  of 2.5 to 3.2 observed on freshwater fishes from Ologe Lagoon, Lagos, Nigeria reported by Kumolu-Johnson and Ndimele (2011).



**Figure 3.** Length-weight relationships of *Labeo coubie*, *Labeo parvus* and *Labeo senegalensis* from the Yapei stretch of the White Volta River. N, Number of fish; W, Weight (grams) and TL, Total Length (cm)



**Figure 4.** Length-weight relationships of *Parailia pellucida*, *Schilbe mystus* and *Distichodus engycephalus* from the Yapei stretch of the White Volta River. N, Number of fish; W, Weight (grams) and TL, Total Length (cm)

The numerical values of  $b$  of the weight-length relationships of *Brycinus leuciscus*, *Hydrocynus forskali*, *Labeo coubie*, *Labeo parvus*, *Labeo senegalensis* and *Schilbe mystus* were within the acceptable and expected range of 2.5 and 3.5 which is typical for tropical fish stocks (Carlander, 1969; Froese, 2006). However, the remaining 3 species: *Odaxothrissa mento*, *Distichodus engycephalus* and *Parailia pellucida* had  $b$  values below 2.5 ( $b < 2.5$ ) and thus indicating negative allometric growth pattern. Additionally on growth pattern, 4 fish species namely *Brycinus leuciscus*, *Labeo senegalensis*, *Labeo parvus* and *Schilbe mystus* were observed to be growing isometrically in weight while 2 species: *Hydrocynus forskali* and *Labeo coubie* had positive allometric growth. Patra *et al.* (2000) stated that the departure of “ $b$ ” values from 3.0 would be due to the seasonal changes while, Le Cren (1951) opined that the deviation of “ $b$ ” value from isometric growth either due to environmental condition or condition of fish. The departure of 5 species from isometric growth in this present study agrees with the findings of the previous authors.

The results on the relative condition factors (Kn) of the 9 species are shown in Table 2. The relative condition factor ranged between 1.00 and 1.09. Subsequently the highest deviation from the expected weight was 0.09 (9%). A t-test of the observed weight ( $W_o$ ) versus the calculated weight ( $W_c$ ) showed no significant differences ( $P > 0.05$ ) between the two weights in all the 9 species.

**Table 1. Descriptive statistics and W-TL relationship parameters for 9 selected fish species of the Yapei Stretch of the White Volta, Ghana (N: sample size; W: weight (g); min: minimum; max: maximum; TL: Total length (cm); S.D.: standard deviation; S.E.: standard error; C.I.: confidence interval; *b*: slope)**

Family/Species	N	Mean W $\pm$ SD (W <sub>min</sub> - W <sub>max</sub> ) grams	Mean TL $\pm$ SD (TL <sub>min</sub> - TL <sub>max</sub> ) cm	W-L Model	Determination Coefficient (r <sup>2</sup> )	<i>b</i>	S.E. of <i>b</i> (95% C.I. of <i>b</i> )	Growth Type
ALESTIDAE <i>Brycinus leuciscus</i>	108	7.51 $\pm$ 1.88 (3.6-11.9)	8.39 $\pm$ 0.69 (6.7-9.9)	W = 0.0118TL <sup>3.026</sup>	0.9004	3.026	0.0978 (2.834-3.218)	Isometric
<i>Hydrocynus forskali</i>	114	100.28 $\pm$ 248.57 (10.80-2040.00)	21.07 $\pm$ 7.9607 (12.50-60.30)	W = 0.0048TL <sup>3.0888</sup>	0.9784	3.0888	0.0434 (3.004-3.174)	Allometric (+)
CLUPEIDAE <i>Odaxothrissa mento</i>	76	7.43 $\pm$ 1.7993 (4.1-10.5)	9.61 $\pm$ 0.8096 (7.1-11.1)	W = 0.0315TL <sup>2.4042</sup>	0.7126	2.4042	0.1775 (2.056-2.752)	Allometric (-)
CYPRINIDAE <i>Labeo coubie</i>	68	226.52 $\pm$ 317.44 (7.50-2160.00)	24.11 $\pm$ 8.50 (9.60-56.20)	W = 0.0064TL <sup>3.1614</sup>	0.941	3.1614	0.0974 (2.970-3.352)	Allometric (+)
<i>Labeo parvus</i>	19	427.10 $\pm$ 340.57 (73.20-910.20)	31.12 $\pm$ 9.98 (17.00-45.00)	W = 0.0103TL <sup>3.0149</sup>	0.9737	3.0149	0.1203 (2.779-3.251)	Isometric
<i>Labeo senegalensis</i>	271	84.32 $\pm$ 80.89 (5.20-620.00)	16.78 $\pm$ 5.29 (8.00-36.40)	W = 0.0193TL <sup>2.8733</sup>	0.808	2.8733	0.0854 (2.706-3.041)	Isometric
DISTICHODONTIDAE <i>Distichodus engycephalus</i>	18	6.94 $\pm$ 1.69 (4.90-10.50)	8.80 $\pm$ 0.88 (7.60-10.6)	W = 0.0489TL <sup>2.2712</sup>	0.9319	2.2712	0.1535 (1.970-2.572)	Allometric (-)
SCHILBEDAE <i>Parailia pellucida</i>	30	4.77 $\pm$ 0.93 (3.10-6.50)	9.23 $\pm$ 0.68 (8.20-10.50)	W = 0.0541TL <sup>2.0045</sup>	0.5764	2.0045	0.3248 (1.368-2.641)	Allometric (-)
<i>Schilbe mystus</i>	37	18.52 $\pm$ 11.34 (5.70-47.90)	14.39 $\pm$ 2.90 (10.00-21.00)	W = 0.0051TL <sup>3.0253</sup>	0.9635	3.0253	0.0995 (2.830-3.220)	Isometric

**Table 2. Mean relative condition factors of the nine (9) selected fish species from the Yapei Stretch of the White Volta, Ghana (N: sample size; Wo: observed weight (g); Wc: calculated weight (g))**

	N	Mean Wo	Mean Wc	Relative Condition Factor Wo/Wc	Mean Deviation (Wo-Wc)/Wc	T-test of Wo Vs. Wc P-values
<i>B. leuciscus</i>	108	7.51	7.50	1.00	0.00	0.99 (P>0.05)
<i>H. forskali</i>	114	100.28	92.63	1.01	0.01	0.79 (P>0.05)
<i>O. mento</i>	76	7.43	7.35	1.01	0.01	0.79 (P>0.05)
<i>L. coubie</i>	68	226.52	221.91	1.03	0.03	0.93 (P>0.05)
<i>L. parvus</i>	19	427.10	424.47	1.01	0.01	0.98 (P>0.05)
<i>L. senegalensis</i>	271	84.32	82.29	1.09	0.09	0.78 (P>0.05)
<i>D. engycephalus</i>	18	6.94	6.92	1.00	0.00	0.97 (P>0.05)
<i>P. pellucida</i>	30	4.77	4.73	1.01	0.01	0.83 (P>0.05)
<i>S. mystus</i>	37	18.52	18.28	1.01	0.01	0.93 (P>0.05)

P>0.05: No significant difference between observed weight (Wo) and calculated weight (Wc)

The relative condition factor being exactly 1.00 for 2 species and very close (1.01) to 1 for 5 species suggested the overall suitability and good state of wellbeing for the fish species.

#### Conclusion

The fitted weight-length power regression models ( $W=aTL^b$ ) generated from this study for the 9 fish species have proven to deliver good relative condition indices are thus suitable for future approximation weights and assessment of the state of wellbeing of the studied fish species.

#### Acknowledgement

The author is grateful to the fishermen of Yapei for their cooperation during the data collection.

#### References

- Abobi, S.M. and Ekau, W. (2013). Length-Weight Relationships and Condition Factors of *Alestes Baremoze*, *Brycinus Nurse* and *Schilbe Intermedius* from the Lower Reaches of White Volta River (Yapei), Ghana. *International Journal of Fisheries and Aquaculture*. 5(6): 152-165
- Bagenal, T.B. and Tesch, F.W. 1978. Age and growth. In: T. Bagenal (ed.), *Methods for assessment of fish production in*

fresh waters, 3rd Edn. IBP Handbook No. 3, Blackwell Science Publications, Oxford:101-136.

3. Bolger, T and Connolly, P.L. (1989). The selection of suitable indices for the measurement and analysis of fish condition. *Journal of Fish Biology* 34:171-182.

4. Braimah, L.I. (2003). Fisheries Management Plan for the Volta Lake, Accra, Ghana. Ministry of Food and Agriculture, Directorate of Fisheries. 77 pp.

5. Carlander, K.D. (1969). Handbook of freshwater fishery biology. Vol. 1. The Iowa State University Press, Ames, IA, pp. 752.

6. Dankwa, H.R., Abban, E.K., Teugels, G.G. (1999). Freshwater fishes of Ghana: identification, distribution, ecological and economic importance. *Anals.Sci. zool.* 283

7. Dulčić, J. and Kraljević, M. (1996). Weight-length relationship for 40 fish species in the Eastern Adriatic (Croatian waters). *Fish. Res.*, 28: 243-251.

8. Enstua-Mensah M, Osei-Abunyew A, Palomares MLD. (1995). Length-weight relationships of fishes from the tributaries of the Volta river, Ghana: Part 1. Analysis of pooled data sets. *Naga, ICLARM Q.* 18, 36-38.

9. FAO: UNDP. (1971). Physico-chemical conditions of Lake Volta, Ghana. Report prepared for the Government of Ghana by FAO of the UN acting as executing agency for the UNDP. Based on the work of C.W. Czernin-Chudenitz. FI: SF:GHA:10: Technical Report. Rome: FAO, 77pp.
10. FAO: UNDP. (1979). Volta Lake research and development project, Ghana. Project findings and recommendations. Terminal Report. Report prepared for the Government of Ghana by FAO of the UN Acting as Executing Agency for UNDP. FAO:FI: DP:GHA: 67:510:71:535. Rome:FAO, 75 pp
11. Froese, R (2006). Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *J. Appl. Ichthyol.*, 22: 241-253.
12. Gayanilo, F.C. and Pauly, D. (1997). The FAO-ICLARM Stock Assessment Tools (FISAT). Reference manual. Rome, FAO, 262p.
13. Gonçalves, J.M.S., Bentes, L., Lino, P.G., Ribeiro, J., Canário, A.V.M. and Erzini, K. (1997). Weight-length relationship for selected fish species of the small-scale demersal fisheries of the south and south-west coast of Portugal. *Fish. Res.*, 30: 253-256.
14. Karna, S. K., Sahoo, D. K. and Panda, S. 2012. Length-Weight Relationship and Growth estimation of *Lates calcarifer* (Bloch) in Chilika lagoon, India. *Bulletin of Environment, Pharmacology & Life Sciences*, 1(3): 61 – 66.
15. King, R.P. (1996). Length-weight relationship of Nigerian Coastal water fishes. *Fishbyte*, 19(4):53-58.
16. Kumolu-Johnson, C.A and Ndimele, P.E. (2011). Length-weight relationships of nine fish species from Ologe Lagoon, Lagos, Nigeria. *African Journal of Biotechnology*. 10 (2): 241-243
17. Kund, G.C., Mishra, G. and Satpathy, D. (2011). Length-weight relationship and relative condition factor of *Mugil cephalus* in Chilika lagoon, east coast of India. *Journal of Applied Zoological Researches*. 22 (2): 153-156.
18. Le Cren, E.D. (1951). The length-weight relationship and seasonal cycles in gonadal weight and condition in perches. *J. Anim. Ecol.* 20: 201-209.
19. McGurk, M.D. (1985). Effects of net capture on the postpreservation morphometry, dry weight, and condition factor of Pacific herring larvae. *Transactions of the American Fisheries Society* 114:348–355.
20. Moutopoulos, D.K. and Stergiou, K.I. (2000). Weight-length and length-length relationships for 40 fish species of the Aegean Sea. *Journal of Applied Ichthyology*, 18:200-203.
21. Ofori-Danson, P.K., Vanderpuye, C.J., de Graaf, G.J. (2001). Growth and mortality of the catfish, *Hemisynodontis membranaceus* (Geoffroy St. Hilaire), in the northern arm of Lake Volta, Ghana. *Fisheries Management and Ecology*. 2001. 8, 37–4.
22. Patra, K.P., Das, S. and Sahu, S. (2000). Seasonal fluctuation of trace metals in *P. Monodon* of Ganjam, East coast of India. *Indian J. fish.* 47 (3): 271-274.
23. Petrakis, G. and Stergiou, K.I. (1995). Weight-length relationship for 33 fish species in Greek waters. *Fisheries Res.*, 21: 465-469.
24. Ricker, W.E. (1975). Computation and interpretation of biological statistics of fish populations. *Bulletin of the Fisheries Research Board of Canada* 191, 382.
25. Santos, M.N., Gaspar, M.B., Vasconcelos, P.V. and Monteiro, C.C. (2002). Weight-length relationship for 50 selected fish species of the Algarve coast (Southern Portugal). *Fisheries Res.*, 9: 289-295.
26. Sokal, R. and Rohlf, F. (1987). *Introduction to Biostatistics*. Freeman, New York, USA.
27. Weatherley, A.H and Gill, H.S. (1987). *The biology of fish growth*. Academic Press, London (UK), 443 p.
28. Wikipedia. (2012). [http://en.wikipedia.org/wiki/Tamale,\\_Ghana](http://en.wikipedia.org/wiki/Tamale,_Ghana) (last accessed: 16.07.2012).